

Comment on the intermediate vortex liquid phase in extreme type-II superconductors

M. V. Feigel'man,

L. D. Landau Institute for Theoretical Physics, 117940 Moscow, Russia

It is shown that disentangled vortex liquid can be a true thermodynamic phase of type-II superconductors in spite of the absence of supercurrent in the direction parallel to the magnetic field.

The problem of the nature of phase transition(s) into a superconductive state in extreme type-II superconductors placed in external magnetic field was recently addressed in a number of numerical simulations¹⁻³ and experiments^{4,5} on YBa₂Cu₃O₇ single crystals. The main issue was formulated long ago⁶ as follows: does an intermediate state of vortex matter ("disentangled vortex liquid", DVL) exist between well-understood Abrikosov lattice and normal metal states? Following early discussions in^{6,7}, the DVL was understood as the state with superconductive current response, $J_z = -\text{const} \cdot \delta A_z$, along the direction of background magnetic field $\mathbf{B} \parallel \hat{z}$. With such an understanding, the results of numerical and experimental studies¹⁻⁵ are apparently conflicting: measurements⁴ and simulations³ indicate the presence of some intermediate vortex liquid state, whereas other measurements⁵ and simulations¹⁻³ demonstrate absence of superconductive response in the vortex liquid. In the present Comment I would like to point out a simple way to reconcile the above observations and to provide some understanding of the vortex state discussed in^{4,3}.

Consider a 3D superconductor with current $\mathbf{J}$ and vector potential $\mathbf{A}$. The longitudinal response of the superconductor in the vortex state is defined by,

$$J_z(\mathbf{q}, q_z) = -P_{zz}(\mathbf{q}, q_z)A_z(\mathbf{q}, q_z), \quad (1)$$

where $\mathbf{q} \perp \hat{z}$. There is a "duality relation" between the longitudinal polarization function of the superconductor P_{zz} and the transverse polarization function $\Pi_\perp(\omega = 0, \mathbf{q})$ of the corresponding 2D Bose-liquid ground-state^{6,7}

$$P_{zz}(\mathbf{q}, q_z = 0) = \frac{1}{\chi^2} \frac{q^2}{q^2 + g^2 \Pi_\perp(\omega = 0, \mathbf{q})}, \quad (2)$$

(in the context of the frustrated XY model², $\Upsilon_z(\mathbf{q}) = P_{zz}(\mathbf{q}, q_z = 0)$ is called the helicity modulus). For the superfluid ground-state of 2D Bose-liquid, $\Pi_\perp(q \rightarrow 0) = \rho_s > 0$, and the duality relation (2) gives a normal longitudinal response $P_{zz} \propto q^2$. For the 2D Bose-liquid ground-state with finite susceptibility χ , like in normal (non-superfluid) liquid, one gets $\Pi_\perp(\mathbf{q} \rightarrow 0) = \chi q^2$, and Eq. (2) gives $\Upsilon_z(q \rightarrow 0) > 0$, i.e. a non-zero helicity modulus. Numerical data¹⁻³ show no evidence of such a liquid state. The point to be made is that these data *do not contradict* to the existence of non-superfluid Bose-liquid ground-state *per se*.

Namely, suppose that a Bose-liquid ground-state possesses a transverse polarization function with an anomalous q -dependence, $\Pi_\perp(\mathbf{q}) \propto |q|^a$ with $0 < a < 2$. The corresponding state of the superconductor will have

$\Upsilon_z(q \rightarrow 0) \propto |q|^{2-a}$, so the $q = 0$ helicity modulus is zero. At the same time, such a ground-state is qualitatively different from the superfluid Bose liquid (i.e. "entangled vortex liquid" in superconductive language); its existence is intrinsically related to the appearance of massless excitation modes with nonzero momentum q , and breakdown of Galilei invariance, as described in⁸. On the other hand, an existence of an intermediate state with such a massless modes is the direct consequence of experimental observations⁴. The only numerical data for the finite- q helicity modulus $\Upsilon(q)$ are provided in², cf. their Fig.3b. The data for $T = 0.54$ seem to be consistent with the above hypothesis, although their accuracy is not yet sufficient to establish this. An additional phase transition above the melting line observed (by counting of vortex lines trajectories) in Ref. 3 can be understood macroscopically as a transition into a state *dual* to the above non-superfluid Bose-liquid.

On a purely theoretical side, arguments in favor of existence of an intermediate nonsuperfluid Bose-liquid ground state were advanced in Ref. 8 where a Bose liquid with a special kind of instantaneous interaction was considered. The peculiar feature of the interaction is that it produces no long-wavelength shear modulus $C_{66}(q \rightarrow 0) = 0$ even in the periodic crystal-like ground state; as a result, the ground state was found to be very fragile. Although a physical realization of the specific model studied in Ref. 8 looks doubtful, it has the same qualitative property, $C_{66}(q \rightarrow 0) = 0$, as the vortex-vortex interaction in the recently studied borocarbide compounds of the RNi₂B₂C family at the critical line $H^*(T)$ of the hexagonal-to-square vortex lattice transition.⁹ We speculate that there may be a fluctuation region around the mean-field transition line $H^*(T)$ between the vortex lattices of different symmetries, and that an anomalous vortex liquid state may occur in borocarbides within this region.

To conclude, a simple way to reconcile apparently conflicting published results¹⁻⁵ on existence of disentangled vortex liquid in HTSC is proposed. I am grateful to V. B. Geshkenbein and A. E. Koshelev for many useful discussions. This research was supported by the RFBR grant #98-02-16252 and the DGA grant #94-1189.

¹ X. Hu *et al* Phys. Rev. Lett. **79**, 3498 (1997).

² P. Olsson and S. Teitel, Phys. Rev. Lett. **82**, 2183, (1999).

- ³ A. K. Nguyen and A. Sudbø, Phys.Rev. B **60**, 15307 (1999).
⁴ A. V. Samoilov *et al*, Phys. Rev. Lett. **76**, 2798 (1996).
⁵ D. Lopez *et al*, Phys. Rev. Lett. **76**, 4034 (1996).
⁶ M. V. Feigel'man *et al*, JETP Lett. **52**, 546 (1990).
⁷ M. V. Feigel'man *et al*, Phys. Rev. B **48**, 16641 (1993).
⁸ M. V. Feigel'man and M. A. Skvortsov, Nucl. Phys. B **506**, 665 (1997).
⁹ U. Yaron *et al*, Nature **382**, 236 (1996); M. R. Eskildsen *et al*, Nature **393**, 242 (1998).